

ATTAKAN C 344 SE

Ina gramu 144 za cypermethrin na 200 za imidacloprid kwa kila lita 1 ya mchanganyiko.

INSECTICIDE / DAWA YA KUUA WADUDU
MFUKO WA KUENDELEZA ZAO LA PAMPA



Pata uhakika kwenye kilimo chako cha pamba



'Kichupa kidogo chenye nguvu kubwa'



Net content : 100 ml
Ujazo : 100 Millilita

Watengenezaji:

Wasambaji:

 **Arysta LifeScience**
ROUTE D ARTIX-B.P. 80
64150 NOGUERES-FRANCE

 **Arysta LifeScience Tanzania Ltd**
Nyerere Road, Plot 127, CSI building-
PO Box 42942 Dar es Salaam, Tanzania
Phone/ fax: (+255)(0) 732 99 18 37

ATTAKAN C 344 SE

Cypermethrin 144 g/l + Imidacloprid 200 g/l, SE

ATTAKAN C 344 SE

Ina gramu 144 za cypermethrin na 200 za imidacloprid kwa kila lita 1 ya mchanganyiko.

ATTAKAN C 344 SE

Is a systemic insecticide with contact and stomach action. It is efficient against bollworms such as Helicoverpa, and against sucking pest

SAFETY PRECAUTIONS

Keep out of children, Keep away from food, water supplies, flammable materials, heat and flame. Wear protective clothing and rubber gloves. Avoid contact with eyes, skin or clothing. Do not eat, drink or smoke in the working area or where the chemical is stored. Avoid breathing vapour or spray mist. Do not apply this product such a manner as to directly expose other persons 10 spray drift. Wash thoroughly with soap and water after handling and before eating or smoking. Immediately remove contaminated clothing and wash before reuse.

FIRST AID

Inhalation: remove from exposure area to fresh air immediately. If breathing has stopped, perform a respiration. Keep person warm and at rest. Treat symptomatically and supportively. Get medical attention immediately.

Eye contact: wash eyes immediately with plenty of water or normal saline, occasionally lifting upper and lower lids, until no evidence of chemical remains (approximately 15- 20 minutes). Get medical attention.

Skin contact: remove contaminated clothing and shoes immediately. Wash affected area with soap or mild detergents and plenty of water until no evidence of chemical remains (approximately 15- 20 minutes). Get medical attention.

Ingestion: remove by gastric lavage and catharsis. Maintain blood pressure and airway. Give oxygen if respiration is depressed. Do not perform gastric lavage if victim is unconscious. Get medical attention immediately.

DIRECTIONS FOR USE

ATTAKAN C 344 SE is a systematic insecticide with contact and stomach action. It gives rapid knock down and a long residual activity efficient against the whole spectrum of bollworms, including Helicoverpa, and against sucking pests such as aphids, white flies and white stainers. **ATTAKAN C344 SE** is applied with a knapsack sprayer or a very low volume sprayer (ULVA)+ at the rate of 100ml/acre.

RE-ENTRY PERIOD: Do not enter treated areas for 4 days unless appropriate protective clothing is worn.

PRE-HARVEST INTERVAL: 3 days.

STORAGE: Store in original container, secured and dried areas, under cool and well ventilated conditions. Prevent cross contamination with other pesticides and fertilizers. Do not store above 38C for extended period of time.

DISPOSAL: Avoid improper disposal of excess concentrate, spray mixture or rinsate. Triple rinse containers puncture and dispose off in an approved sanitary landfill. Do not reuse containers.

ADVICE FOR PHYSICIANS: No specific antidote. Treat symptomatically and supportively.

ATTAKAN C 344 SE

Ni kiuwa dudu kinachopenyeza ndani ya mmea kuuwa wadudu kama sumu ya tumbo au mguso. Inadhibiti wadudu aina ya viwavi, funza na wadudu wanaoifyonza.

TAHADHARI

Hifadhi mbali na watoto, vyakula vya binadamu na mifugo, vitu vinavyowaka kwa urahisi, joto, moto pamoja na vyanzo vya maji. Vaa mavazi ya kinga na glavu za mpira. Epuka dawa isingie machoni, isiguse ngozi au mavazi. Epuka kuvuta mvuke au mnyunyizio wa dawa. Usile, usinywe wala kuvuta katika eneo la kazi au mahali dawa hii inapohifadhiwa. Usitumie dawa hii kwa njia inayoweza kusababisha madhara kwa binadamu. Nawa vizuri kwa maji na sabuni baada ya kutumia dawa hii. Vua nguo za kazi mara tu baada ya kazi na hakikisha zimefuliwa kabla ya kuzitumia tena.

HUDUMA YA KWANZA

Ikivutwa: Mhamishe haraka aliyedhurika na kumpeleka kwenye hewa safi. Kama mgonjwa anashindwa kupumua, apatiwe oksijeni, hakikisha anawekwa katika hali ya joto na apumzike. Tibu kulingana na dalili na kwa msaada tu. Mpeleke kwa daktari.

Ikiingia machoni: safisha macho haraka kwa maji mengi huku ukiyafumbua mara kwa mara mpaka utakapohakikisha hakuna dalili zozote za dawa. (Dakika zisizopungua 15-20). Mpeleke kwa daktari.

Ikigusa ngozi: vua nguo zilizolowa dawa pamoja na viatu. Safisha sehemu iliyoathirika kwa sabuni na maji mengi kwa muda wa dakika 15-20. Mpeleke aliyedhurika kwa daktari ikimezwa.

MAELEZO YA MATUMIZI

ATTAKAN 344 SE ni kiuwa dudu kinachopenyeza ndani ya mmea kuuwa wadudu kama sumu ya tumbo au mguso. Inafanya kazi haraka kuangusha wadudu na inaedelea kufanya kazi kwa muda mrefu baadaye. Inadhibiti viwavi wote ikijumulisha na wadudu wanaoifyonza kama chawa, nzi weupe, vidukari na kadhalika.

ATTAKAN C 344 SE hutumika kwa kipimo cha milimita 100 kwa hekta na kunyunyizwa kwa kutumia mabomba ya kawaida au yale ya betri (ulva+)

KUINGIA ENEO LILILONYUNYIZIWA

Usiingie eneo lililonyunyiziwa kwa muda wa siku 4 bila kuvaa mavazi ya kinga

USALAMA WA MAZAO: Subiri siku 3 baada ya kunyunyiza ndipo uvune ili kuepuka madhara.

KUHIFADHI: Hifadhi dawa hii kwenye vyombo vyake vya awali vilivyofungwa madhubuti, sehemu yenye usalama, kavu, isiyo na joto na yenye kuruhusu hewa ya kutosha. Zuia isichanganyikane na dawa nyingine pamoja na mbolea. Usihifadhi kwa muda mrefu kwenye sehemu yenye joto zaidi ya nyuzi 38 C.

KUTEKETEZA MABAKI YA DAWA NA VYOMBO VITUPU

Epuka kumwaga ovyo mabaki ya dawa na misuuzo. Suza vyombo vilivyokwisha dawa mara tatu, halafu vitoboe na kuvitupa kwenye jalala lililoidhinishwa..

USHAURI KWADAKTARI

Hakuna antidoti maalum. Kwa msaada tu tibu kulingana na dalili

